Theory and Practice of Science and Technology
2026, VOL. 7, NO. 1, 83-86
DOI: 10.47297/taposatWSP2633-456917.20260701

Soil Organic Carbon Distribution and Its Influencing Factors in Mountainous Regions

Weihuan Li, Dong Yang, Yingying Zhou

Northwest Normal University, Lanzhou, Gansu, 730070, China

ABSTRACT
Soil which is organic as a matter of fact and has carbon (which is called SOC) plays a really important part in the cycle of carbon at a global level. This part has an effect on the fertility belonging to the soil, the productivity of the ecosystem,and the act of regulating the climate. Regions which are of a mountainous kind, and these regions are characterized by a topography that is complex and micro - climates that show diversity, serve as important yet changeable storage places for SOC. To gain an understanding about the spatial distribution of SOC in these specific areas and the factors that control it is very important for the accurate calculation of carbon and the sustainable management of land. This paper conducts an examination on the main patterns associated with the distribution of SOC in mountainous terrain and performs an analysis of the primary factors that have an impact, including the climate, the vegetation,the form of the land, the features of the soil, and the actions made by humans. The combination shows that the storage of soil organic carbon is very uneven in mountain areas. Frequently, it shows an increase while the height goes up until it reaches a certain limit due to the changes in temperature and amount of rain. Nevertheless, this pattern is adjusted by the local situations. The interaction among these factors leads to complex changes of soil organic carbon. Strategies that are useful for the improvement of the storage of soil organic carbon in areas that are mountain - covered must take into consideration this control system that has many aspects. They should combine the protection of the ecological surroundings with practices of using the land that are suitable so as to decrease the effects of climate change.

KEYWORDS
Soil organic carbon; Mountainous ecosystems; Spatial distribution; Influencing factors; Carbon sequestration

1 Introduction

Soil organic carbon forms a major portion of the carbon pool on the land, and it holds approximately two to three times more carbon when compared to the atmosphere. The alterations of it are extremely significant for the well - being of the soil, the long - term viability of agriculture, and the global climate system [1]. Ecosystems in mountain regions cover around one - quarter of the Earth's land area and are considered as areas that are responsive and vital for the circulation of carbon. In the areas of mountains, there exists a kind of interaction which is complex among the steep gradients belonging to the environment, the various geological substrates, and the different types related to land cover. This particular interaction results in a distribution of the soil organic carbon (SOC) that shows a characteristic of being heterogeneous in the aspect of space and being stratified in the vertical direction. This sort of heterogeneity brings forward challenges for the process of the assessment of carbon stocks on a large - scale basis. However, it also offers a natural place for people to carry out the study of the factors that have control over the formation, the stabilization, and the loss of the SOC [2].

It is necessary to precisely carry out the action of quantifying the stocks of soil organic carbon (SOC) and gain an understanding of the mechanisms which control them in mountain - like areas. This action gives support to the efforts of making a prediction regarding the response of terrestrial carbon sinks to climate change. It provides information for the policies that are formulated for the conservation of the ecosystem and offers guidance for the practices that are carried out for the purpose of mitigating climate change by means of enhancing carbon sequestration.Even if a great quantity of studies have been done in various mountain areas all over the world, the action of putting together the general rules that control the distribution of soil organic carbon (SOC) still stays as a main goal in the realm of science. This paper has the purpose of performing a review and having a talk about the general patterns of the distribution of SOC in areas that are mountain - like and methodically doing an analysis of the factors that are of natural origin and those that are brought about by human actions and that have an impact on these patterns, with an emphasis on providing insights that are related for the management of the ecological system without making a mention of particular locations.

2 Patterns of Soil Organic Carbon Distribution in Mountainous Regions

The distribution of SOC in mountains is non-uniform, exhibiting distinct horizontal and vertical patterns influenced by the dramatic environmental changes over short distances.

2.1 Vertical Distribution with Soil Depth

The concentration of soil organic carbon (SOC) typically decreases in an exponential manner as the depth of the soil gets greater. The layer located on the surface (0 - 20 cm), and this layer is filled with the litter from recent plants and the exudates of roots, has the highest density of SOC. This layer which is active plays a very important role in the cycling of nutrients in the short - term. In the layers that are beneath the surface and at greater depths, the SOC consists of materials that are more difficult to decompose and have undergone humification, as well as carbon that has been stabilized and is associated with the surfaces of minerals. In the mountainous areas, the depth of the soil itself is frequently restricted by the gradient of the slope and the process of erosion, and this erosion process can cut off the profiles of the soil and limit the potential for the total storage of SOC [3].

2.2 Horizontal and Elevational Patterns

The pattern which people have studied the most is the variation which changes along elevational gradients. A trend which is widespread and has been noticed in many mountains which are moist, mild - climate, and tropical is that there is a starting sort of rise in stocks of soil organic carbon (SOC) as the elevation rises. This situation mainly results from the reduction which occurs in temperature, which causes the rates of microbial decomposition to decline, and frequently there is an increase in rainfall and dampness.which causes an event of bringing about a boosting of the productivity of plants to a greater extent and an event of adding litter[4]. The happening of the occurrence of the peak of SOC storage often takes place under the conditions that are cold and wet in the forest belts at mid - elevation. After reaching a particular elevation, when going into alpine meadows or tundra zones, low temperatures seriously restrict the growth of plants and the net primary production, and this leads to a decrease in the inputs of SOC, and thus there are lower total amounts even though the decomposition process is slow. This results in the creation of a pattern of SOC storage that has a hump - shaped appearance along the elevation gradient.

Speaking from the side direction, the distribution of the carbon which is organic in the soil (SOC) is seriously influenced by the position of the topography. Landforms which are sunken, for example, depressions, valleys, and foot - slopes, have a likelihood to gather SOC due to the transportation of water, sediments, and the carbon which is organic and dissolved down the slope. This causes the creation of the deposits of soil which are thicker and a content of moisture which is higher. In contrast,Land - forms which have a convex shape, such as ridges and upper slopes that are steep, are often shown by the occurrence of erosion, soils which are thinner, moisture levels that are lower, and because of this, a reduction in the process of storing SOC. The aspect, which refers to the direction of the slope, also exerts an influence. Slopes that are in the shade and face towards the north (in the Northern Hemisphere) are usually cooler and wetter. This kind of situation is beneficial for the build - up of SOC when making a comparison with slopes that face towards the south, which are warmer and drier.

3 Influencing Factors of Soil Organic Carbon Distribution

The distribution patterns described above are the result of the integrated effects of multiple biotic and abiotic factors.

3.1 Climate Factors

Climate acts as a driver which is basic both at regional level and global level. Temperature shows an opposite relation with the amount of SOC in most of the biomes. Temperatures that are lower at elevations that are higher have an effect that inhibits the activity of microbes and enzymes. This leads to a slowing down in the breakdown of organic matter and brings about its gathering.Precipitation exerts an effect on Soil Organic Carbon (SOC) by exerting an impact on both the productivity of plants (input) and the rates of decomposition. Adequate moisture plays a part in promoting the growth of plants, which results in an increase in the fall of litter and the biomass of roots. However, excessive moisture can cause waterlogging and anaerobic conditions, which slow down the decomposition process but may also impose restrictions on certain types of vegetation. The interaction between temperature and precipitation serves to determine the hydrothermal conditions that, fundamentally, exercise control over the balance between the inputs of SOC and the outputs of SOC.

3.2 Vegetation and Land Cover

Vegetation type stands as a main source for soil organic carbon (SOC) by means of the production of litter, The turnover happening to roots, along with the deposition taking place of rhizodeposits. Communities belonging to different types of vegetation make contributions that are of varying amounts and qualities of organic matter. For instance, forests consisting of coniferous trees produce litter which has a high content of lignin and decomposes at a slow speed,

and this has the potential to result in the formation of thick organic horizons [5]. Forests made up of broadleaf trees may go through a faster turnover of litter but possess a larger biomass of roots. Grasslands make a substantial contribution to SOC via systems that are dense and fine of roots. The change occurring in land - use, such as the conversion that happens of natural forests into cropland or pasture, is one of the most remarkable anthropogenic factors that cause the loss of SOC, and it disturbs the equilibrium existing between input and decomposition.

3.3 Topographic Factors

Topography causes the making of modifications to the climate together with ecological processes at local scales, and it makes the creation of micro - environments. Elevation acts as a replacement for the changes in temperature and precipitation. Slope gradient along with slope position exert an influence on soil erosion, deposition, and the redistribution of water. Slopes which are more gentle and landscape positions which are lower are good for the carrying out of the accumulation of SOC. Aspect has an impact on the receiving of solar radiation, soil temperature, and evapotranspiration, and by doing this action, it affects the activity of microbes and the composition of communities of plants. The curvature state of the slope (regardless of whether it is convex or concave) decides whether an area is a net source or a net sink in relation to water and sediments, and this directly has an impact on SOC.

3.4 Soil Properties

Characteristics that the soil inherently has play a role in determining the ability which is to stabilize and also to protect the soil organic carbon (SOC). It is critical that the texture of the soil exists；soils having a fine texture (including clay and silt) possess a surface area that is larger and are able to form complexes of organo - mineral which can protect the SOC from the attack of microbes both physically and chemically. Mineralogy, especially the situation where reactive iron and aluminum oxides are present, has an effect of enhancing the stabilization of SOC. The density of the soil bulk has an impact on the calculation of the stocks of SOC and is in itself affected by the content of organic matter and the way of management. The pH has an influence on the composition and the activity of the microbial community, and as a result, it affects the pathways of decomposition. The depth of the soil and the material of the parent set up the ultimate volume in physical terms and the context in mineralogical terms for the storage of SOC.

3.5 Anthropogenic Activities

Activities that are carried out by human beings are having an impact that is increasingly great on the dynamics of soil organic carbon (SOC) which is inside mountainous regions. Practices that are related to agricultural activities, for example the activity of tillage, cause an acceleration in the process of decomposition and the occurrence of erosion of SOC. On the other hand, the farming way of building terraces can play a function in helping to reduce erosion and maintain the SOC in an intact state.The intensity degree of grazing has the ability to cause the soil to turn into a compacted state, bring about a decrease in the vegetation cover, and lead to the occurrence of changes in the plant community structure, which very frequently leads to the exhaustion of SOC. The actions of cutting down forests and the actions of planting new forests or the actions of re - establishing forests have extremely profound and opposite impacts on the SOC stocks. The act of applying fertilizers can result in an increase in the biomass production and the input of SOC.yet it might also bring about some changes to the rates of the process of decomposition. The practices which are related to sustainable land management, like the farming way that does not involve tilling the land, the act of combining agriculture and forestry, and the practice of carrying out controlled grazing, are very much necessary with the purpose of keeping or making better the SOC in mountain landscapes that are affected by humans.

4 Implications for Carbon Sequestration and Management

The soil organic carbon (SOC) in mountains has a heterogeneous nature, and this indicates that strategies for the purpose of management and conservation need to be made specific to the particular site. At the same time, it is necessary to take into account the interaction of local factors. Areas that are of high priority for the act of conserving SOC are generally those that possess high existing stocks (for example, forests located at mid - elevation, wetlands, stable footslopes) and have a high susceptibility to the occurrence of loss (for example, steep slopes under intensive use). Practices that lead to a decrease in soil erosion are beneficial in a general sense. This is because erosion not only removes SOC but also causes a disturbance to the soil structure.

Making an enhancement to carbon sequestration calls for practices that bring an increase to carbon inputs and/or bring about a reduction in decomposition losses. This involves promoting the recovery of natural vegetation, putting into

practice sustainable forestry, carrying out the restoration of degraded lands with species that have deep roots, and adopting agricultural techniques that are friendly to the soil on mountain farmlands. Policy frameworks ought to give encouragement to such practices by means of incentives and integrate the protection of SOC into broader goals related to the conservation of biodiversity, the protection of water resources, and the adaptation to climate change in mountainous regions.

5 Conclusion

Mountain areas are components that are dynamic and important of the carbon pool of the global soil. The way of distribution of soil organic carbon (SOC) inside them is marked by obvious heterogeneity in space that is driven by gradients which are vertical (depth) and horizontal (elevation, topography). This kind of distribution is controlled by a set that is complex of factors which interact with each other: climate sets the constraints that are broad, vegetation offers the input that is primary, topography redistributes resources and creates niches that are local, properties of soil decide the capacity of stabilization, and activities of humans bring in changes that are dynamic. Understanding this hierarchy of controls is very important for developing models of prediction that are accurate and policies of land management that are effective. Research in the future should focus on doing a better job of quantifying stocks of SOC in mountain areas where sampling is insufficient, making clear the mechanisms of stabilization of SOC under conditions that are different, and evaluating the effectiveness in the long term of various interventions of management for the purpose of carbon sequestration. Protecting and increasing SOC in ecosystems that are mountainous is not just a strategy for climate mitigation but also a basis for their stability that is ecological and the well - being of communities that rely on their resources.

References

[1] CHEN Y, AMANI-BENI M, ZHANG R, et al. Evolution of population distribution and its influencing factors in the poverty-stricken mountainous region of Southwest China from 2000 to 2020 [J]. Palgrave Communications, 2024, 10(1): 156.

[2] CHEN Y, AMANI-BENI M, ZHANG R, et al. Evolution of population distribution and its influencing factors in the poverty-stricken mountainous region of Southwest China from 2000 to 2020 [J]. Humanities & Social Sciences Communications, 2024, 11(1): 167.

[3] WANG L, FAN H, WU Z H, et al. Spatial pattern and influencing factors of topsoil organic carbon contents in hilly and mountainous regions of Eastern Sichuan in China [J]. Transactions of the Chinese Society of Agricultural Engineering, 2024, 40(23): 169-178.

[4] ZHANG L, XU T, BAI Y, et al. Historical and future dynamics of soil organic carbon and driving mechanisms in mountainous soils of China [J]. Catena, 2024, 244: 108456.

[5] Xinlei C , Ying C . Distribution Characteristics and Influence Factors of Key Rural Tourism Villages: A Case Study of Beijing[J]. Journal of Landscape Research, 2025, 17(3):33-39.

[6] Gao J , Zhang Y , Zuo L .The optimal explanatory power of soil erosion and water yield in karst mountainous areas[J].Journal of Geographical Sciences, 2023, 33(10):2077-2093.